TURKISH JOURNAL OF SCIENTIFIC RESEARCH

Volume I Issue III ISSN:2148-5135



VITAMINS

Asatullayev Rustamjon Bakhtiyarovich
Scientific supervisor:
Student: **Mamadaliyeva Malika**

ARTICLE INFO:

Received:08.11.2026
Revised: 09.11.2026
Accepted:10.11.2026

KEY WORDS:

vitamins, human health, nutrition, fat-soluble vitamins, water-soluble vitamins, metabolism, immune system, vitamin deficiency, dietary sources, nutritional balance.

ABSTRACT

Vitamins are essential organic compounds required in small amounts for the normal growth, development, and functioning of the human body. They play a significant role in metabolism, immune system regulation, tissue repair, bone formation, and the maintenance of healthy skin, vision, and nervous system function. Although most vitamins cannot be synthesized by the human body in sufficient quantities, they can be obtained through a balanced diet and, when necessary, dietary supplements. Vitamins are classified into two main groups: fat-soluble vitamins, including vitamins A, D, E, and K, and water-soluble vitamins, including vitamin C and the B-complex vitamins. Each vitamin performs specific physiological functions, and deficiencies may lead to various health problems. This article examines the classification, biological functions, dietary sources, and clinical significance of vitamins. It also highlights the importance of adequate vitamin intake and the potential risks associated with excessive supplementation. Understanding the role of vitamins is essential for promoting nutritional health, preventing deficiency-related disorders, and supporting overall well-being.

Vitamins are essential micronutrients that support numerous biological processes in the human body. Although they are required in relatively small quantities compared with proteins, carbohydrates, and fats, they are indispensable for maintaining normal physiological functions. Vitamins contribute to energy metabolism, cellular growth, tissue development, immune responses, and the regulation of various biochemical reactions. Since the body cannot produce

most vitamins in sufficient amounts, an adequate supply through food is necessary to maintain health and prevent nutritional deficiencies.

Vitamins are generally divided into two categories according to their solubility: fat-soluble and water-soluble vitamins. Fat-soluble vitamins include vitamins A, D, E, and K. These vitamins can be absorbed along with dietary fats and stored in the liver and fatty tissues. Water-soluble vitamins include vitamin C and the eight B vitamins, which are involved in numerous metabolic processes. Most water-soluble vitamins are not stored extensively in the body, although vitamin B12 is an important exception because significant amounts can be stored in the liver. Consequently, regular dietary intake is particularly important for maintaining adequate levels of many water-soluble vitamins.

Vitamin A is essential for normal vision, immune function, cellular differentiation, and the maintenance of healthy skin and mucous membranes. It can be obtained from animal-based foods, such as eggs, dairy products, and liver, as well as from plant foods containing carotenoids, including carrots, pumpkins, and dark green leafy vegetables. Vitamin A deficiency may cause night blindness and, in severe cases, more serious visual impairment. However, excessive intake of preformed vitamin A, particularly through high-dose supplements, can be toxic.

Vitamin D plays an important role in calcium and phosphorus absorption, bone mineralization, and muscle function. The human body can produce vitamin D in the skin following exposure to ultraviolet B radiation, while dietary sources include oily fish, egg yolks, and fortified foods. Inadequate vitamin D levels can contribute to impaired bone mineralization, causing rickets in children and osteomalacia in adults. Vitamin D is also being studied for its wider effects on immune and metabolic functions, although not all proposed benefits of supplementation are firmly established.

Vitamin E functions primarily as an antioxidant, helping protect cell membranes from oxidative damage. It is found in vegetable oils, nuts, seeds, and certain green vegetables. Although vitamin E deficiency is uncommon in healthy individuals, it can occur in people with conditions that impair fat absorption and may lead to neurological and muscular problems. Vitamin K is necessary for normal blood clotting and also contributes to bone metabolism. Important dietary sources include leafy green vegetables, such as spinach, kale, and broccoli. Severe vitamin K deficiency can increase the risk of abnormal bleeding.

The B-complex vitamins are essential for energy metabolism, red blood cell formation, and normal nervous system function. Vitamin B1, or thiamine, helps the body metabolize carbohydrates and supports nerve function. Vitamin B2, or riboflavin, participates in energy-producing reactions, while vitamin B3, or niacin, contributes to cellular metabolism and DNA maintenance. Vitamin B5, known as pantothenic acid, is required for the synthesis of coenzyme A, which is involved in numerous metabolic pathways. Vitamin B6 supports amino acid metabolism, neurotransmitter synthesis, and hemoglobin formation. Biotin, or vitamin B7, participates in reactions involved in the metabolism of fats, carbohydrates, and proteins.

Folate, or vitamin B9, is particularly important for DNA synthesis, cell division, and the formation of red blood cells. Adequate folate intake before conception and during early pregnancy reduces the risk of neural tube defects in developing fetuses. Vitamin B12, or cobalamin, is essential for neurological function, DNA synthesis, and red blood cell production. It is naturally found primarily in animal-derived foods, including meat, fish, eggs, and dairy products, although fortified foods and supplements can provide alternatives. Vitamin B12 deficiency may cause megaloblastic anemia and neurological complications. Individuals following strict vegan diets may require a reliable fortified food or supplement source of this vitamin.

Vitamin C, also known as ascorbic acid, is a water-soluble vitamin that contributes to collagen synthesis, wound healing, antioxidant protection, and immune system function. It also enhances the absorption of non-heme iron from plant-based foods. Citrus fruits, berries, kiwi, peppers, tomatoes, and broccoli are valuable dietary sources. Severe vitamin C deficiency causes scurvy, a condition associated with bleeding gums, poor wound healing, fatigue, and connective tissue abnormalities. Since vitamin C is sensitive to heat and can be lost during prolonged cooking, consuming a variety of fresh or minimally cooked fruits and vegetables can help preserve its content.

Vitamin deficiencies can develop because of inadequate dietary intake, malabsorption, certain medical conditions, increased nutritional requirements, or the use of particular medications. The symptoms vary according to the vitamin involved and may include fatigue, anemia, impaired vision, weakened bones, skin changes, and neurological problems. Some deficiencies develop gradually and may remain undetected for a long time. Therefore, appropriate clinical assessment and laboratory testing may be necessary when a deficiency is suspected.

A balanced and varied diet is generally the most effective way for healthy individuals to obtain the vitamins they need. Fruits, vegetables, whole grains, legumes, nuts, seeds, dairy products or suitable alternatives, eggs, fish, and other nutrient-rich foods can collectively provide a broad range of essential vitamins. Dietary supplements may be beneficial in specific circumstances, including pregnancy, certain dietary restrictions, diagnosed deficiencies, or medical conditions that interfere with nutrient absorption. However, supplements should not automatically be considered a replacement for a balanced diet. Excessive intake of some vitamins, particularly fat-soluble vitamins, can cause harmful effects, and high doses of certain water-soluble vitamins can also produce adverse reactions. Supplementation should therefore be guided by individual needs and appropriate professional advice.

In conclusion, vitamins are indispensable micronutrients that contribute to the normal functioning of nearly every major body system. Their roles include supporting metabolism, strengthening physiological defenses, maintaining healthy bones and tissues, and enabling normal blood formation and neurological activity. Both inadequate and excessive vitamin intake can negatively affect health, making nutritional balance essential. A varied diet, appropriate

supplementation when necessary, and greater awareness of nutritional requirements can help prevent vitamin-related disorders and promote long-term health and well-being.

References

1. World Health Organization, & Food and Agriculture Organization of the United Nations. (2004). Vitamin and mineral requirements in human nutrition (2nd ed.). World Health Organization. https://iris.who.int/handle/10665/42716
2. National Institutes of Health, Office of Dietary Supplements. (n.d.). Dietary supplement fact sheets. https://ods.od.nih.gov/factsheets/list-all/
3. Gropper, S. S., & Smith, J. L. (2021). Advanced Nutrition and Human Metabolism (8th ed.). Cengage Learning.
4. Mahan, L. K., & Raymond, J. L. (Eds.). (2020). Krause and Mahan's Food & the Nutrition Care Process (15th ed.). Elsevier.
5. Combs, G. F., Jr., & McClung, J. P. (2017). The Vitamins: Fundamental Aspects in Nutrition and Health (5th ed.). Academic Press.
6. Kennedy, D. O. (2016). B vitamins and the brain: Mechanisms, dose and efficacy—A review. Nutrients, 8(2), Article 68. https://doi.org/10.3390/nu8020068
7. Carr, A. C., & Maggini, S. (2017). Vitamin C and immune function. Nutrients, 9(11), Article 1211. https://doi.org/10.3390/nu9111211
8. National Institutes of Health, Office of Dietary Supplements. (n.d.). Vitamin D: Fact sheet for health professionals. https://ods.od.nih.gov/factsheets/VitaminD-HealthProfessional/